# **Elabscience**®


### APC Anti-Mouse CD28 Antibody[37.51]

Catalog Number: E-AB-F1026UE

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Mouse
Host	Syrian Hamster
Isotype	Syrian Hamster IgG
Clone No.	37.51
Isotype Control	[Product E-AB-F09763E]
Conjugation	APC
Conjugation Information	APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the
	reagent to obtain optimal results [The recommended concentration is 0.1-1 $\mu$ g/10 <sup>6</sup> cells in 100 $\mu$ L volume].

Data



C57BL/6 murine splenocytes are stained with FITC Anti-Mouse CD3 Antibody and APC Anti-Mouse CD28 Antibody (Left). Splenocytes are stained with FITC Anti-Mouse CD3 Antibody and APC Syrian Hamster IgG Isotype Control (Right).

Preparation & Storage	
Storage	Keep as concentrated solution.
Shipping	This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze. Ice bag
Antigen Information	
Alternate Names	CD28;Cd28;T-cell-specific surface glycoprotein CD28
Uniprot ID	P31041

### For Research Use Only

## **Elabscience**®

Gene ID Background

#### 12487

CD28 is a 44 kD glycoprotein, also known as Tp44 or T44. It is a member of the lg superfamily, expressed on thymocytes, most peripheral T cells, and NK cells. In association with CD80 (B7-1) and CD86 (B7-2), CD28 acts as the second signal for T and NK cell activation and proliferation. The 37.51 antibody has been reported to augment in vitro T cell proliferation and cytokine production, and promote CTL development.

For Research Use Only