

Purified Anti-Mouse CD185 Antibody[L138D7]

catalog number: AN008940P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Mouse
Immunogen	Recombinant Mouse CD185 protein
Host	Rat
Isotype	Rat IgG2b, κ
Clone	L138D7
Purification	>98%, Protein A/G purified
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

Applications

Recommended Dilution

FCM	2 $\mu\text{g}/\text{mL}$ (0.5×10^6 - 1×10^6 cells)
------------	--

Data



C57/BL6 Mouse splenocytes were stained with 0.2 μg Purified Anti-Mouse CD185 Antibody[L138D7](Right) and 0.2 μg Rat IgG2b, κ Isotype Control (Left), followed by APC-conjugated Goat Anti-Rat IgG Secondary Antibody, then anti-Mouse B220 FITC-conjugated Monoclonal Antibody.

Preparation & Storage

Storage	Store at 4°C valid for 12 months or -20°C valid for long term storage, avoid freeze / thaw cycles.
Shipping	Ice bag

Background

CD185 is also known as CXCR5. It is the receptor for chemokine CXCL13/BLC, which is chemotactic for B cells. CXCR5 is expressed on B cells and a subset of T cells in the spleen, neuronal tissue, lymph nodes, and bone marrow. It is important for migration of B cells into the B cell follicles of the spleen and Peyer's patches. Follicular helper T cells (Tfh) also express CXCR5 and the ability of these cells to migrate to the lymph node is modulated by the balanced expression of CCR7 and CXCR5.

For Research Use Only