

## Recombinant Human STIM1/GOK Protein (His Tag)

**Catalog Number:** PKSH031120

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human STIM1/GOK protein Met 1-Asp 213, with an C-terminal His
<b>Calculated MW</b>	23.3 kDa
<b>Observed MW</b>	33-38 kDa
<b>Accession</b>	NP_003147.2
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 97 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 97 % as determined by reducing SDS-PAGE.

### Background

#### For Research Use Only

Stromal interaction molecule 1, also known as STIM1 and GOK, is a cell membrane, a single-pass type I membrane protein and an endoplasmic reticulum membrane protein. STIM1 / GOK is ubiquitously expressed in various human primary cells and tumor cell lines. It contains one EF-hand domain and one SAM (sterile alpha motif) domain. STIM1 / GOK plays a role in mediating Ca<sup>2+</sup> influx following depletion of intracellular Ca<sup>2+</sup> stores. It acts as a Ca<sup>2+</sup> sensor in the endoplasmic reticulum via its EF-hand domain. Upon Ca<sup>2+</sup> depletion, STIM1 / GOK translocates from the endoplasmic reticulum to the plasma membrane where it activates the Ca<sup>2+</sup> release-activated Ca<sup>2+</sup> (CRAC) channel subunit, TMEM142A / ORAI1. Transfection of STIM1 / GOK into cells derived from a rhabdoid tumor and from a rhabdomyosarcoma that do not express detectable levels of STIM1 can induce cell death, suggesting a possible role in the control of rhabdomyosarcomas and rhabdoid tumors. Defects in STIM1 are the cause of immune dysfunction with T-cell inactivation due to calcium entry defect type 2 (IDTICED2) which is an immune disorder characterized by recurrent infections, impaired T-cell activation and proliferative response, decreased T-cell production of cytokines, lymphadenopathy, and normal lymphocyte counts and serum immunoglobulin levels.

## For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017