

A Reliable Research Partner in Life Science and Medicine

FITC Anti-Human CD8 Antibody[SK1]

Catalog Number: AN00337C

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human;Rhesus;Cynomolgus

Host Mouse

Isotype Mouse IgG1, κ

Clone No. SK1

Isotype Control FITC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792C]

Conjugation FITC

Conjugation Information FITC is designed to be excited by the Blue laser (488 nm) and detected using an optical

filter centered near 530 nm (e.g., a 525/40 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Data



Staining of normal human peripheral blood cells with FITC Anti-Human CD8 Antibody[SK1] (filled gray histogram) or FITC Mouse IgG1, κ Isotype Control (empty black histogram). Cells in the lymphocytes gate were used for analysis.

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

 Alternate Names
 T8;Leu2

 Uniprot ID
 P01732

 Gene ID
 925

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience®

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Background

CD8a is a 32-34 kD type I glycoprotein. It forms a homodimer (CD8a/a) or heterodimer (CD8a/b) with CD8b. CD8, also known as T8 and Leu2, is a member of the immunoglobulin superfamily found on the majority of thymocytes, a subset of peripheral blood T cells, and NK cells (which express almost exclusively CD8a homodimers). CD8 acts as a co-receptor with MHC class I-restricted T cell receptors in antigen recognition and T cell activation and has been shown to play a role in thymic differentiation. Two domains in CD8a are important for function: the extracellular IgSF domain binds the $\alpha3$ domain of MHC class I and the cytoplasmic CXCP motif binds the tyrosine kinase p56 Lck

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web:www.elabscience.com
 Email:techsupport@elabscience.com