Recombinant Mouse Interleukin-13/IL-13 protein (His Tag)

Catalog Number: PDMM100037



Note: Centrifuge before opening to ensure complete recovery of vial contents.

-				
	A C	Cri	nti	nn
\mathbf{L}	\mathbf{c}	cri		$\mathbf{v}_{\mathbf{H}}$

 Species
 Mouse

 Mol_Mass
 14.3 kDa

 Accession
 P20109

Bio-activity Not validated for activity

Properties

Purity > 95% as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU/mg of the protein as determined by the LAL method

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5%

Mannitol.

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution of

0.5 mg/mL. Concentration is measured by UV-Vis.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Mouse interleukin 13 (mIL-13) is a pleiotropic cytokine produced by activated Th2 cells. IL-13 induces B cell proliferation and immunoglobin production. It contains a four helical bundle with two internal disulfide bonds. Mouse IL13 shares 5 8% sequence identity with human protein and exhibits cross-species activity. IL13 signals via receptor IL13R (type2, IL4R) and activates STAT-6. IL13 initially binds IL-13Rα1 with low affinity and triggers association of IL4Rα, generating a high affinity heterodimeric receptor IL13R and eliciting downstream signals. IL13 also binds IL-13Rα2 with high affinity, which plays a role in a negative feedback system of IL13 signaling. IL13 is an important mediator of allergic inflammation and disease.

For Research Use Only