

A Reliable Research Partner in Life Science and Medicine

APC Anti-Human CD183/CXCR3 Antibody[G025H7]

Catalog Number: E-AB-F1156E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Host Mouse

 Isotype
 Mouse IgG1, κ

 Clone No.
 G025H7

Isotype Control APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792E]

Conjugation APC

Conjugation Information APC is designed to be excited by the Red (627-640 nm) laser and detected using an

optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.


Applications Recommended usage

FCM Each lot of this antibody is quality control tested by flow cytometric analysis. The amount

of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for

individual use.

Data



Human peripheral blood lymphocytes are stained with PE Anti-Human CD3 Antibody and APC Anti-Human CD183/CXCR3 Antibody (Left). Lymphocytes are stained with PE Anti-Human CD3 Antibody and APC Mouse IgG1, κ Isotype Control (Right).

Preparation & Storage

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping lce bag

Antigen Information

Alternate Names CKR-L2;CXC-R3;CXCR-3;CXCR3;GPR9

 Uniprot ID
 P49682

 Gene ID
 2833

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web:www.elabscience.com Email:techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

Background

Human CXCR3, also known as GPR9, is a chemokine receptor that binds CXCL9, CXCL10, and CXCL11. It is a 38 kD seven-pass transmembrane receptor coupled to G-protein. CXCR3 is highly expressed by T cells (Th1), natural killer cells (NK cells), dendritic cells, mast cells, alveolar macrophages, eosinophils, and human airway epithelial cells. CXCR3 is important for effector lymphocyte recruitment into inflamed tissue in various inflammatory and autoimmune diseases, such as chronically inflamed liver, Crohn's disease, rheumatoid arthritis, multiple sclerosis, and inflammatory skin diseases.

Fax: 1-832-243-6017