A Reliable Research Partner in Life Science and Medicine

# Recombinant Human CDK2AP2 Protein (E.coli, His Tag)

Catalog Number: PKSH030818

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Species Human

Source E.coli-derived Human CDK2AP2 protein Met 1-Thr 126, with an C-terminal His

 Calculated MW
 14.5 kDa

 Observed MW
 18&12 kDa

 Accession
 O75956

**Bio-activity** Not validated for activity

#### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** Please contact us for more information.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, 15% glycerol, pH 7.5


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

## Data



> 95 % as determined by reducing SDS-PAGE.

### Background

CDK2AP2 belongs to the CDK2AP family. Members of this family of proteins are cell-growth suppressors; associating with and influencing the biological activities of important cell cycle regulators in the S phase including monomeric non-phosphorylated cyclin-dependent kinase 2 (CDK2) and DNA polymerase alpha/primase. CDK2AP2 contains 5 distinct g t-ag introns. Transcription produces 7 different mRNAs; 6 alternatively spliced variants and 1 unspliced form. There are 2 non overlapping alternative last exons and 4 validated alternative polyadenylation sites. The mRNAs appear to differ splicing versus retention of 3 introns. CDK2AP2 plays a role in regulating self-renewal of mouse embryonic stem cells ( mESC) under permissive conditions; and cell survival during differentiation of the mESC into terminally differentiated cell types.

#### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017