

Recombinant Mouse Syndecan-4/SDC4 Protein (Fc Tag)

Catalog Number: PKSM040519

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Mouse
Source	HEK293 Cells-derived Mouse Syndecan-4/SDC4 protein Met1-Val146, with an C-terminal hFc
Calculated MW	40.5 kDa
Observed MW	33 kDa
Accession	O35988
Bio-activity	Immobilized human MDK at 10 µg/ml (100 µl/well) can bind mouse SDC4-Fc with a linear range of 0.16-1.25 µg/ml.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

SDC4 (Syndecan-4), also known as Syn4, is a transmembrane heparan sulfate proteoglycan that co-operates with integrins during cell-matrix interactions for the assembly of focal adhesions and actin stress fibers and in the phosphorylation of focal adhesion kinase (FAK) on Tyr397. Syndecan-4 plays roles in the formation of focal adhesions and stress fibers. The cytoplasmic domain of syndecan-4 interacts with a number of signalling and structural proteins, and both extracellular and cytoplasmic domains are necessary for regulated activation of associated transmembrane receptors. Syndecan-4/SDC4 is a heparan sulfate proteoglycan and works as a coreceptor for various growth factors. SDC4 deficiency limits neointimal formation after vascular injury by regulating vascular smooth muscle cells (VSMCs) proliferation and vascular progenitor cells (VPCs) mobilization. Therefore, SDC4 may be a novel therapeutic target for preventing arterial restenosis after angioplasty.