

## Recombinant Mouse Syndecan-4/SDC4 Protein (Fc Tag)

**Catalog Number:** PKSM040519

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Mouse
<b>Source</b>	HEK293 Cells-derived Mouse Syndecan-4/SDC4 protein Met1-Val146, with an C-terminal hFc
<b>Calculated MW</b>	40.5 kDa
<b>Observed MW</b>	33 kDa
<b>Accession</b>	O35988
<b>Bio-activity</b>	Immobilized human MDK at 10 µg/ml (100 µl/well) can bind mouse SDC4-Fc with a linear range of 0.16-1.25 µg/ml.

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

#### For Research Use Only

SDC4 (Syndecan-4), also known as Syn4, is a transmembrane heparan sulfate proteoglycan that co-operates with integrins during cell-matrix interactions for the assembly of focal adhesions and actin stress fibers and in the phosphorylation of focal adhesion kinase (FAK) on Tyr397. Syndecan-4 plays roles in the formation of focal adhesions and stress fibers. The cytoplasmic domain of syndecan-4 interacts with a number of signalling and structural proteins, and both extracellular and cytoplasmic domains are necessary for regulated activation of associated transmembrane receptors. Syndecan-4/SDC4 is a heparan sulfate proteoglycan and works as a coreceptor for various growth factors. SDC4 deficiency limits neointimal formation after vascular injury by regulating vascular smooth muscle cells (VSMCs) proliferation and vascular progenitor cells (VPCs) mobilization. Therefore, SDC4 may be a novel therapeutic target for preventing arterial restenosis after angioplasty.

## For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017