Elabscience®

Biotin Anti-Mouse CD274/PD-L1 Antibody[10F.9G2]

Catalog Number: E-AB-F1132B

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Mouse
Host	Rat
lsotype	Rat IgG2b, κ
Clone No.	10F.9G2
Isotype Control	Biotin Rat IgG2b, κ Isotype Control[LTF-2] [Product E-AB-F09843B]
Conjugation	Biotin
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.
Applications	Recommended usage
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. For flow
	cytometric staining, the suggested use of this reagent is \leq 1.0 µg per 10 ⁶ cells in 100 µL volume or 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Preparation & Storage	
Storage	Keep as concentrated solution.
	This product can be stored at 2-8°C for 12 months. Do not freeze.
Shipping	Ice bag
Antigen Information	
Alternate Names	B7H1PD-L1PDCD1L1PDCD1LG1PDL1;B7 homolog 1;B7-H;B7-H1;Programmed cell death ligand 1
Uniprot ID	Q9EP73
Gene ID	60533
Background	CD274, also known as B7-H1 or programmed death ligand 1 (PD-L1), is a 40 kD type I transmembrane protein and a member of the B7 family within the immunoglobulin receptor superfamily. It is expressed on T cells, B cells, NK cells, dendritic cells, IFN-γ activated endothelial cells, and monocytes. B7-H1 is one of the ligands of PD-1. The interaction of B7-H1 with PD-1 plays an important role in the inhibition of T cell responses. Other studies have shown that B7-H1 is able to costimulate T cell growth and cytokine production. CD274 is involved in costimulation essential for T cell proliferation and production of IL-10 and IFN-γ, in an IL-2-dependent and a PD-1-independent manner. Its interaction with PD-1 inhibits T cell proliferation and cytokine production.