

Mouse IgG Antibody Pair Set

Catalog No. E-KAB-0078

Applications

ELISA

Synonyms IgG

Kit components & Storage

Title	Specifications	Storage
Mouse IgG Capture Antibody	1 vial, 100 µg	Store at -20°C for one year. Avoid freeze / thaw cycles.
Mouse IgG Detection Antibody (Biotin)	1 vial, 50 µL	Store at -20°C for one year. Avoid freeze / thaw cycles.

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Product Information

Items		Characteristic (E-KAB-0078)	
		Mouse IgG Capture Antibody	Mouse IgG Detection Antibody (Biotin)
Immunogen Information	Immunogen	Native Protein	Native Protein
	Swissprot	P01863	
Product details	Reactivity	Mouse	Mouse
	Host	Goat	Goat
	Conjugation	Unconjugated	Biotin
	Concentration	0.5mg/mL	/
	Buffer	PBS with 0.04% Proclin 300, 50% glycerol, pH 7.4	PBS with 0.04% Proclin 300, 1% protective protein, 50% glycerol, pH 7.4
	Purify	Antigen Affinity	Antigen Affinity
	Specificity	Detects Mouse IgG in ELISAs.	

For Research Use Only

Applications

Mouse IgG Sandwich ELISA Assay:

	Recommended Concentration/Dilution	Reagent	Images																
ELISA Capture	0.5-4μg/mL	Mouse IgG Capture Antibody	 <table><caption>Approximate data points from the standard curve</caption><thead><tr><th>Mouse IgG concentration (ng/mL)</th><th>Optical Density</th></tr></thead><tbody><tr><td>1</td><td>0.05</td></tr><tr><td>2</td><td>0.1</td></tr><tr><td>5</td><td>0.2</td></tr><tr><td>10</td><td>0.4</td></tr><tr><td>20</td><td>0.8</td></tr><tr><td>50</td><td>1.5</td></tr><tr><td>100</td><td>2.5</td></tr></tbody></table>	Mouse IgG concentration (ng/mL)	Optical Density	1	0.05	2	0.1	5	0.2	10	0.4	20	0.8	50	1.5	100	2.5
Mouse IgG concentration (ng/mL)	Optical Density																		
1	0.05																		
2	0.1																		
5	0.2																		
10	0.4																		
20	0.8																		
50	1.5																		
100	2.5																		
ELISA Detection	1:1000-1:10000	Mouse IgG Detection Antibody (Biotin)																	

Note: This standard curve is only for demonstration purposes. A standard curve should be generated for each assay!

Background

Constant region of immunoglobulin heavy chains. Immunoglobulins, also known as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into immunoglobulins-secreting plasma cells. Secreted immunoglobulins mediate the effector phase of humoral immunity, which results in the elimination of bound antigens. The antigen binding site is formed by the variable domain of one heavy chain, together with that of its associated light chain. Thus, each immunoglobulin has two antigen binding sites with remarkable affinity for a particular antigen. The variable domains are assembled by a process called V-(D)-J rearrangement and can then be subjected to somatic hypermutations which, after exposure to antigen and selection, allow affinity maturation for a particular antigen.

For Research Use Only