## **Elabscience**®

## **BMP5** Polyclonal Antibody

catalog number: E-AB-17735

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description  |  |  |
|--|--|--|
| Reactivity   | Human;Mouse  |  |
| Immunogen  | Synthetic peptide of human BMP5  |  |
| Host   | Rabbit   |  |
| Isotype  | IgG  |  |
| Purification   | Antigen affinity purification  |  |
| Conjugation  | Unconjugated   |  |
| Buffer   | Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol. |  |
| Applications   | Recommended Dilution   |  |
| WB   | 1:500-1:2000   |  |
| Data   |  |  |
|  | KDa 1 2   130  |  |
| Western blot analysis of Human placenta tissue and Human |  |  |

right lower lung tissue lysates using BMP5 Polyclonal

Antibody at dilution of 1:450

## **Observed-MW:Refer to figures**

Calculated-MW:52 kDa

| Preparation & Storage |   |
|-----------------------|---|
| Storage               | Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.                 |
| Shipping              | The product is shipped with ice pack, upon receipt, store it immediately at the |
|                       | temperature recommended.  |

## Background

Bone morphogenetic protein-5 (BMP5), is a member of the transforming growth factor-beta (TGF-beta) superfamily. The superfamily includes large families of growth and differentiation factors. Bone morphogenetic proteins were originally identified by an ability of demineralized bone extract to induce endochondral osteogenesis in vivo in an extraskeletal sit e. These proteins are synthesized as prepropeptides, cleaved, and then processed into dimeric proteins. BMP5 may act as an important signaling molecule within the trabecular meshwork and optic nerve head, and may play a potential role in

glaucoma pathogenesis. BMP5 is differentially regulated in multiple human cancers.