

## ITCH Polyclonal Antibody

**catalog number: E-AB-91779**

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Reactivity</b>	Human;Mouse;Rat
<b>Immunogen</b>	Recombinant fusion protein of human ITCH
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Purification</b>	Affinity purification
<b>Buffer</b>	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

### Applications


<b>WB</b>	1:500-1:2000
<b>IHC</b>	1:50-1:100

### Data





Western blot analysis of extracts of 293T cells using ITCH Polyclonal Antibody at 1:500 dilution.

**Observed-MV:100 kDa**  
**Calculated-MV:103 kDa**




Immunohistochemistry of paraffin-embedded rat brain using ITCH Polyclonal Antibody at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.




Western blot analysis of extracts of various cell lines using ITCH Polyclonal Antibody at dilution.

**Observed-MV:100 kDa**  
**Calculated-MV:103 kDa**



Immunohistochemistry of paraffin-embedded human lung cancer using ITCH Polyclonal Antibody at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



### For Research Use Only

Immunohistochemistry of paraffin-embedded human placenta using ITCH Polyclonal Antibody at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.	Immunohistochemistry of paraffin-embedded mouse brain using ITCH Polyclonal Antibody at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.
--	---

## Preparation & Storage

<b>Storage</b>	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
<b>Shipping</b>	The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

## Background

This gene encodes a member of the Nedd4 family of HECT domain E3 ubiquitin ligases. HECT domain E3 ubiquitin ligases transfer ubiquitin from E2 ubiquitin-conjugating enzymes to protein substrates, thus targeting specific proteins for lysosomal degradation. The encoded protein plays a role in multiple cellular processes including erythroid and lymphoid cell differentiation and the regulation of immune responses. Mutations in this gene are a cause of syndromic multisystem autoimmune disease. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.

## For Research Use Only