

**IFNAR2 (C-6His), Human, Recombinant****Cat. No. : GPCK162****产品信息**

物种	Human
表达宿主	Human Cells
序列信息	Ile27-Lys243
检索号	P48551
标签	C-6His
分子量	25.8 kDa
有效期	12 months

**产品特性**

内毒素 (EU/μg)	< 0.1
保存	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
运输	Ambient temperature or ice pack.
制剂	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
复溶	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

**背景介绍**

Interferon  $\alpha/\beta$  Receptor 2 (IFN- $\alpha/\beta$  R2) is a single-pass type I membrane Protein which belongs to the type II Cytokine Receptor family. It complexes with IFN- $\alpha/\beta$  R1 to form the signaling Receptor complex for the family of  $\alpha$  and  $\beta$  IFN subtypes. By alternative splicing, IFN- $\alpha/\beta$  R2 can exist as a secreted soluble Protein or as a type I membrane Protein. IFN- $\alpha/\beta$  R2 is the principal Ligand binding subunit of the Receptor. Ligand binding is stabilized by the subsequent association with IFN- $\alpha/\beta$  R1, resulting in the formation of a signaling ternary Receptor complex. IFNAR2 was detected in most lymphocytes, monocytes, and granulocytes, although IFNAR2 expression was higher in the monocytes and granulocytes than in the lymphocytes. Among the lymphocyte subsets, IFNAR2 showed high expression in natural killer (NK) cells and low expression in T lymphocytes. Isoform 1 and isoform 3 of IFNAR2 are directly involved in signal transduction due to their interaction with the TYR kinase, JAK1. Isoform 1 also interacts with the transcriptional factors, STAT1 and STAT2. Both forms are potent inhibitors of type I IFN activity.

