

A Reliable Research Partner in Life Science and Medicine

# APC Anti-Human CD70 Antibody[113-16]

Catalog Number: AN00424E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Reactivity Human Host Mouse

**Isotype** Mouse  $\lg G1$ ,  $\kappa$ 

**Clone No.** 113-16

Isotype Control APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792E]

**Conjugation** APC

**Conjugation Information** APC is designed to be excited by the Red (627-640 nm) laser and detected using an

optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.

### Applications Recommended usage

**FCM** 

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

### **Data**



Staining of the U266 cell line with APC Anti-Human CD70 Antibody[113-16] (filled gray histogram) or APC Mouse IgG1,  $\kappa$  Isotype Control (empty black histogram). Total viable cells were used for analysis.

## **Preparation & Storage**

**Storage** Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping lce bag

### **Antigen Information**

Alternate Names CD27 ligand;CD27L;Ki-24 antigen;Tumor necrosis factor ligand superfamily;member 7

Web: www.elabscience.cn

(TNFSF7)

 Uniprot ID
 P32970

 Gene ID
 970

# For Research Use Only



# **Elabscience Biotechnology Co., Ltd.**

A Reliable Research Partner in Life Science and Medicine

## **Background**

CD70, also known as CD27L, is a 50 kD type II transmembrane glycoprotein and member of the tumor necrosis factor superfamily. CD70 is expressed on activated T, B and NK cells, activated plasmacytoid dendritic cells (pDCs), and chronic B cell lymphocytic leukemia and large B cell lymphomas. CD70 costimulates T cell proliferation and differentiation. It plays a role in the pDC-induced B cell differentiation. The ligand of CD70 is CD27.