

JNK1/2/3 Polyclonal Antibody

catalog number: E-AB-70345

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity	Mouse;Rat
Immunogen	KLH conjugated Synthetic peptide corresponding to human JNK1
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer, 1% protein protectant and 50% glycerol.


Applications

Recommended Dilution


IHC

1:200-1:500


Data



Immunohistochemistry analysis of paraffin-embedded Mouse lung using JNK1/2/3 Polyclonal Antibody at dilution of 1:200.



Immunohistochemistry analysis of paraffin-embedded Mouse brain using JNK1/2/3 Polyclonal Antibody at dilution of 1:200.



Immunohistochemistry analysis of paraffin-embedded Rat brain using JNK1/2/3 Polyclonal Antibody at dilution of 1:200.

Preparation & Storage

Storage

Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping

The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

Background

For Research Use Only

c-Jun N-terminal kinases (JNKs) phosphorylate and augment transcriptional activity of c-Jun. JNKs originate from three genes that yield ten isoforms through alternative mRNA splicing, including JNK1 α 1, JNK1 β 1, JNK2 α 1, JNK2 β 1 and JNK3 α 1, which represent the p46 isoforms, and JNK1 α 2, JNK1 β 2, JNK2 α 2, JNK2 β 2 and JNK3 β 2, which represent the p54 isoforms. JNKs coordinate cell responses to stress and influence regulation of cell growth and transformation. The human JNK1 (PRKM8, SAPK1, MAPK8) gene maps to chromosome 10q11.22 and shares 83% amino acid identity with JNK2. JNK1 is necessary for normal activation and differentiation of CD4 helper T (TH) cells into TH1 and TH2 effector cells. Capsaicin activates JNK1 and p38 in Ras-transformed human breast epithelial cells. Nitrogen oxides (NOx) upregulate JNK1 in addition to c-Fos, c-Jun and other signaling kinases, including MEKK1 and p38.