

Recombinant Human CD136/MST1R Protein (His Tag)



Catalog Number:PKSH031069

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	CD136;CDw136;PTK8;RON
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Leu 571
Accession	Q04912
Calculated Molecular Weight	60.0 kDa
Observed molecular weight	70 kDa
Tag	C-His

Properties

Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

The tyrosine kinase receptor, macrophage-stimulating 1 receptor (MST1R), a c-met-related tyrosine kinase, also known as the Ron receptor or CD136, controls cell survival and motility programs related to invasive growth. As tyrosine kinase receptor comprised of an extra-cellular domain, MST1R protein contains the ligand binding pocket and an intracellular region where the kinase domain is located. MST1R signaling may be involved in the regulation of macrophage and T-lymphocyte activation in vivo during injury. This assessment of gene expression indicates the importance of genetic factors in contributing to lung injury, and points to strategies for intervention in the progression of inflammatory diseases. It had been shown that MST1R/CD136 plays a critical role in Ni-induced lung injury in mice. The overexpression of MSP, MT-SP1, and MST1R was a strong independent indicator of both metastasis and death in human breast cancer

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human CD136/MST1R Protein (His Tag)



Catalog Number:PKSH031069

patients and significantly increased the accuracy of an existing gene expression signature for poor prognosis. Stimulation of MST1R leads to its transphosphorylation and the ultimate activation of numerous intracellular signaling pathways, such as the classical mitogen-activated protein kinase pathway, the phosphatidylinositol (PI)3-kinase pathway, and the JNK pathway.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com