



A Reliable Research Partner in Life Science and Medicine

## PE/Cyanine 5.5 Anti-Mouse CD49b/pan-NK cells Antibody [DX5]

Catalog Number: E-AB-F1116UI

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Mouse Rat Host Isotype Rat IgM, ĸ Clone No. DX5

[Product E-AB-F09773I] Isotype Control

PE/Cyanine 5.5 Conjugation

**Conjugation Information** PE/Cyanine5.5 is designed to be excited by the Blue (488 nm), Green (532 nm) and

yellow-green (561 nm) lasers and detected using an optical filter centered near 690 nm

(e.g., a 690/50 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

**Applications** Recommended usage

**FCM** Each lot of this antibody is quality control tested by flow cytometric analysis. Please

> check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use. We suggest each investigator should titrate the reagent to obtain optimal results [The recommended concentration is 0.1-1 µg/10<sup>6</sup> cells

in 100 µL volume].

**Preparation & Storage** 

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

**Antigen Information** 

**Alternate Names** CD49 antigen-like family member B;CD49b;Collagen receptor;GPla;Integrin alpha-2;

Platelet membrane glycoprotein la;VLA-2 subunit alpha;pan-NK cells

**Uniprot ID** Q62469 Gene ID 16398

**Background** DX5 antigen has been recently characterized as CD49b. It is a 150 kD integrin α chain

> also known as α2 integrin, VLA-2 α chain, and integrin α2 chain. CD49b non-covalently associates with CD29 (β1 integrin) to form the CD49b/CD29 complex known as VLA-2, a receptor for collagen and laminin. CD49b is expressed on platelets, the majority of NK cells, NKT cells, and a small subset of CD8+ T cells (this population can be significantly increased following viral infection). DX5 is used for the identification and isolation of NK cells, and is especially useful for identifying NK cells in mice lacking the

NK1.1 antigen.

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Web:www.elabscience.com

Email:techsupport@elabscience.com