

Recombinant CSF-1R Monoclonal Antibody

catalog number: AN301500L

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human;Mouse
Immunogen	Recombinant human CSF-1R fragment
Host	Rabbit
Isotype	IgG, κ
Clone	A195
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.


Applications Recommended Dilution

IHC	1:500-1:2000
------------	--------------


Data



Immunohistochemistry of paraffin-embedded Human colon using CSF-1R Monoclonal Antibody at dilution of 1:2000.



Immunohistochemistry of paraffin-embedded Human tonsil using CSF-1R Monoclonal Antibody at dilution of 1:1000.



Immunohistochemistry of paraffin-embedded Mouse colon using CSF-1R Monoclonal Antibody at dilution of 1:1000.

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

Tyrosine-protein kinase that acts as cell-surface receptor for CSF1 and IL34 and plays an essential role in the regulation of survival, proliferation and differentiation of hematopoietic precursor cells, especially mononuclear phagocytes, such as macrophages and monocytes. Promotes the release of proinflammatory chemokines in response to IL34 and CSF1, and thereby plays an important role in innate immunity and in inflammatory processes. Plays an important role in the regulation of osteoclast proliferation and differentiation, the regulation of bone resorption, and is required for normal bone and tooth development. Required for normal male and female fertility, and for normal development of milk ducts and acinar structures in the mammary gland during pregnancy. Promotes reorganization of the actin cytoskeleton, regulates formation of membrane ruffles, cell adhesion and cell migration, and promotes cancer cell invasion.

For Research Use Only