## **Elabscience**®

### **TTR Polyclonal Antibody**

#### catalog number: E-AB-18211

Note: Centrifuge before opening to ensure complete recovery of vial contents.


1:50-1:200


Description	
Reactivity	Human;Mouse;Rat
Immunogen	Fusion protein of human TTR
Host	Rabbit
Isotype	IgG
Purification	Antigen affinity purification
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.
Applications	Recommended Dilution
WB	1:500-1:2000

### IHC

#### Data

WB





cancer tissue using TTR Polyclonal Antibody at dilution of


1:50(×200)

Western blot analysis of Mouse serum using TTR Polyclonal Immunohistochemistry of paraffin-embedded Human liver

Antibody at dilution of 1:600

#### **Observed-MW:Refer to figures**

#### Calculated-MW:16 kDa



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using TTR Polyclonal Antibody at dilution of  $1:50(\times 200)$ 

Preparation & Storage	
Storage Shipping	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

#### Background

#### For Research Use Only

# **Elabscience**®

Transthyretin (TTR) is a plasma transport protein for thyroxine and retinol,through the association with retinol-binding protein. It is a homotetrameric protein synthesized mainly in liver,choroid plexus,retinal pigment epithelium,and pancrea s.Within the CNS,TTR is the only known protein synthesized solely by the choroid plexus. Mutant and wildtype TTR give rise to various forms of amyloid deposition (amyloidosis). Defects in TTR are the cause of amyloidosis transthyretin-related (AMYL-TTR),hyperthyroxinemia dystransthyretinemic euthyroidal (HTDE) and carpal tunnel syndrome type 1 (CTS1). In addition,positive immunostaining for TTR has been reported as a sensitive diagnostic marker of choroid plexus tumors. (22103483)

For Research Use Only