HA-Tag Polyclonal Antibody

catalog number: **E-AB-40523**


Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description

Reactivity	All
Immunogen	Synthetic peptide corresponding to HA tag conjugated to keyhole limpet haemocyanin.
Host	Rabbit
Isotype	Rabbit IgG
Purification	Antigen Affinity Purification
Buffer	Phosphate-buffered solution, pH 7.2, containing 0.05% non-protein stabilizer. Dialyze to completely remove the stabilizer prior to labeling.

Applications	Recommended Dilution
WB	1:5000-1:10000
IP	3μg/sample
IF	1:800-1600
FCM	0.2μg


Data



Western blot with Anti HA-Tag Polyclonal Antibody at dilution of 1:10000. Lane 1:293T Ttrasfected with HA-Tag cell lysate,Lane 2:293T cell lysate.

Observed-MW:28 kDa


Calculated-MW:28 kDa



Western blot with Anti HA-Tag Polyclonal Antibody at dilution of 1:10000. Lane 1:293T Ttrasfected with HA-Tag cell lysate,Lane 2:293T cell lysate.

Observed-MW:28 kDa

Calculated-MW:28 kDa



Western blot with Anti HA-Tag Polyclonal Antibody at dilution of 1:10000. Lane 1:293T Ttrasfected with HA-Tag cell lysate,Lane 2:293T cell lysate.

Observed-MW:28 kDa

Calculated-MW:28 kDa



Western blot with Anti HA-Tag Polyclonal Antibody at dilution of 1:10000. Lane 1:293T Ttrasfected with HA-Tag cell lysate,Lane 2:293T cell lysate.

Observed-MW:28 kDa

Calculated-MW:28 kDa

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.

Shipping	The product is shipped with ice pack,upon receipt,store it immediately at the temperature recommended.

Background

The HA tag is derived from the HA-molecule corresponding to amino acids 98-106. It has been extensively used as a general epitope tag in expression vectors. Many recombinant proteins have been engineered to express the HA tag, which does not appear to int

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com