

## A Reliable Research Partner in Life Science and Medicine

# **Caspase-7 Polyclonal Antibody**

catalog number: E-AB-60327

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

Reactivity Human; Mouse; Rat

Immunogen Recombinant fusion protein of human Caspase-7 (NP 001218.1).


Host Rabbit Isotype IgG

Purification Affinity purification


Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

#### **Recommended Dilution Applications** WB 1:500-1:2000 IHC 1:50-1:200 IF 1:50-1:200

### Data




Caspase-7 Polyclonal Antibody at dilution of 1:1000.




Western blot analysis of extracts of various cell lines using Immunohistochemistry of paraffin-embedded Rat heart using Caspase-7 Polyclonal Antibody at dilution of 1:100 (40x lens).

## Observed-MV:37 kDa Calculated-MV:28 kDa/31 kDa/34 kDa/37 kDa



Immunohistochemistry of paraffin-embedded Human lung cancer using Caspase-7 Polyclonal Antibody at dilution of 1:100 (40x lens).



Immunofluorescence analysis of HeLa cells using Caspase-7 Polyclonal Antibody

## Preparation & Storage

Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Storage

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

## Background

### For Research Use Only

Toll-free: 1-888-852-8623 Web:www.elabscience.com

## **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

This gene encodes a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. The precursor of the encoded protein is cleaved by caspase 3 and 10, is activated upon cell death stimuli and induces apoptosis. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.

Fax: 1-832-243-6017