Recombinant Human TGFBI/BIGH3 Protein (His Tag)

Catalog Number: PKSH031520

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Species	Human
Source	HEK293 Cells-derived Human TGFBI/BIGH3 protein Met 1-His 683, with an C-terminal
	His
Calculated MW	74.0 kDa
Observed MW	65 kDa
Accession	NP_000349.1
Bio-activity	Not validated for activity
Properties	
Purity	>75 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants
	before lyophilization.
	Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	
	KDa M
	116
	66.2
	45.0
	35.0
	25.0
	18.4 14.4

> 75 % as determined by reducing SDS-PAGE.

Background

TGFBI is an RGD-containing protein that binds to type I, II and IV collagens. The RGD motif is found in many extracellular matrix proteins modulating cell adhesion and serves as a ligand recognition sequence for several integrins. TGFBI plays a role in cell-collagen interactions and may be involved in endochondrial bone formation in cartilage. TGFBI is induced by transforming growth factor-beta and acts to inhibit cell adhesion. Mutations in TGFBI are associated with multiple types of corneal dystrophy. TGFBI can bind to type I, II, and IV collagens. This adhesion protein may play an important role in cell-collagen interactions. In cartilage, TGFBI may be involved in endochondral bone formation. Loss of the TGFBI is sufficient to induce specific resistance.

For Research Use Only