

Recombinant Human CD32b/FCGR2B protein (His tag)

Catalog Number:PDMH100113



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms	Low Affinity Immunoglobulin Gamma Fc Region Receptor II-b;IgG Fc Receptor II-b;CDw32;Fc-Gamma RII-b;Fc-Gamma-RIIb;FcRII-b;CD32;FCGR2B;FCG2;IGFR2
Species	Human
Expression Host	HEK293 Cells
Sequence	Thr43-Pro217
Accession	P31994
Calculated Molecular Weight	19.1 kDa
Observed molecular weight	30 kDa
Tag	C-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Background

Fc γ RIIB is a low affinity receptor that recognizes the Fc portion of IgG. The human CD32 group consists of Fc γ RIIA, Fc γ RIIB, and Fc γ RIIC proteins that share 94-99% sequence identity in their extracellular domains but differ substantially in their transmembrane and cytoplasmic domains. Fc γ RII protein is expressed on cells of both myeloid and lymphoid lineages as well as on cells of non-hematopoietic origin. Fc γ RIIB has an intrinsic cytoplasmic immunoreceptor tyrosine-based inhibitory motif (ITIM) and delivers an inhibitory signal upon ligand binding. Ligation of Fc γ RIIB on B cells down-regulates antibody production and in some circumstances may promote apoptosis. Co-ligation of Fc γ RIIB on dendritic cells inhibits maturation and blocks cell activation. Fc γ RIIB may also be a target for monoclonal antibody therapy for malignancies. Fc γ RIIB plays an important negative-regulating role through modulating the signals from activation receptors.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017