## Recombinant Mouse LAIR1 Protein (aa 22-141, His Tag)

Catalog Number: PKSM041099



Description		
Species	Mouse	
Mol_Mass	14.4 kDa	
Accession	Q8BG84	
Bio-activity	Not validated for activity	
Properties		
Purity	> 95 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80	
	°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of	
	reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.	
	Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants	
	before lyophilization.	
	Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	

Note: Centrifuge before opening to ensure complete recovery of vial contents.

kDa 120 90 60 40	MK	R
30		a sence
Elabsolenie 20		Sec. 1
14		-

> 95 % as determined by reducing SDS-PAGE.

## Background

Data

Leukocyte-associated Ig-like receptor-1 (LAIR-1) is an inhibitory receptor of the Ig superfamily that is structurally related to inhibitory members of KIR and ILT/CD85 families. It is expressed on immune cells, including NK cells, T cells, B cells, monocytes, immature neutrophils, dendritic cells and most thymocytes. The 253 amino acid (aa) type I transmembrane (TM) protein contains a 21 aa signal sequence, a 124 aa extracellular domain (ECD), a 20 aa TM domain and a 98 aa cytoplasmic domain. The ECD includes one C2-type Ig-like domain and two potential N-linked glycosylation sites. Tyrosine phosphorylation of two cytoplasmic ITIM motifs results in recruitment of phosphatases and down-regulation of signaling through activating receptors. LAIR1 shows high-affinity binding of collagens that results in inhibition of degranulation in a basophilic leukemia cell line.

## For Research Use Only