Elabscience®


APC Anti-Human CD147 Antibody[HIM6]

Catalog Number: E-AB-F1056E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description				
Reactivity	Human			
Host	Mouse			
lsotype	Mouse IgG1, κ			
Clone No.	HIM6			
Isotype Control	APC Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792E]			
Conjugation	APC			
Conjugation Information	APC is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).			
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% sodium azide and 1% BSA.			
Applications	Recommended usage			
FCM	Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 μ L of antibody per test (million cells in 100 μ L staining volume or per 100 μ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.			

Data



Human peripheral blood lymphocytes are stained with APC Anti-Human CD147 Antibody (filled gray histogram). Unstained lymphocytes (empty black histogram) are used as control.

Preparation & Storage		
Storage	Keep as concentrated solution.	
	This product can be stored at 2-8°C for 12 months. Please protected from prolonged exposure to light and do not freeze.	
Shipping	Ice bag	
Antigen Information		
Alternate Names	5F7;BSG;Basigin;CD147;Collagenase stimulatory factor;EMMPRIN;Extracellular matrix metalloproteinase inducer;Leukocyte activation antigen M6;OK blood group antigen; TCSF;Tumor cell-derived collagenase stimulatory factor	
Uniprot ID	P35613	

For	Resea	rch U	Jse C)nly
-----	-------	-------	-------	------

Elabscience®

Gene ID
Background

682

CD147, also known as neurothelin or basigin, is a member of the lg superfamily. It is a 55-65 kD type I transmembrane glycoprotein which is primarily expressed on leukocytes, erythrocytes, platelets, and endothelial cells. CD147 is reported to have a function during embryonal brain development and/or play a role in integrin-mediated adhesion in brain endothelia.