## Recombinant Rat IgG2A Fc/Igg-2a Protein

Catalog Number: PKSR030448



| Description    |  |  |
|----------------|--|--|
| Species        | Rat  |  |
| Mol_Mass       | 35.2 kDa   |  |
| Accession      | P20760   |  |
| Bio-activity   | Not validated for activity   |  |
| Properties     |  |  |
| Purity         | > 95 % as determined by reducing SDS-PAGE.   |  |
| Endotoxin      | < 1.0 EU per µg of the protein as determined by the LAL method.                          |  |
| Storage        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 |  |
|                | °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of      |  |
|                | reconstituted samples are stable at $< -20^{\circ}$ C for 3 months.                      |  |
| Shipping       | This product is provided as lyophilized powder which is shipped with ice packs.          |  |
| Formulation    | Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, pH 8.5.                    |  |
|                | Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants         |  |
|                | before lyophilization.   |  |
|                | Please refer to the specific buffer information in the printed manual.                   |  |
| Reconstitution | Please refer to the printed manual for detailed information.                             |  |
| Data           |  |  |

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| kDa       | МК   | R |
|-----------|------|---|
| 120<br>90 |      |   |
| 60        |      |   |
| 40        | -    |   |
| 30        | -    | - |
| 20        | head |   |
|           |      |   |
| 14        |      |   |

> 95 % as determined by reducing SDS-PAGE.

## Background

Immunoglobulin G (IgG) is a type of antibody composed of four peptide chains—two identical heavy chains and two identical light chains arranged in a Y-shape typical of antibody monomers. There are four IgG subclasses (IgG1, 2, 3, and 4) in humans, named in order of their abundance in serum. The IgG2a isotype was able to interact very efficiently with FcgammaR, and share approximately 64-78% amino acid sequence identity with IgG1 and IgG2b in the heavy chain constant region domains, CH1, CH2 and CH3, which cause the diffrence in the interaction with complement, NK and mast cells, induction or inhibition by specific cytokines, and ability to cross the placenta.

## For Research Use Only