Elabscience®

XBP1 Polyclonal Antibody

catalog number: E-AB-68333

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description			
Reactivity	Human;Mouse;Rat		
Immunogen	Recombinant fusion prote	Recombinant fusion protein of human XBP1	
Host	Rabbit		
Isotype	IgG		
Purification	Affinity purification		
Buffer	Phosphate buffered solut	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.	
Applications	Recommended Dilu	Recommended Dilution	
WB	1:500-1:2000		
Data			
۳۵۵۵ ۵۵۵۵ ۵۵۵۵ ۵۶۵۵ ۵۶۵۵ ۵۶۵۶ ۵۶۵۶ ۵۶۵۶	racts of various cell lines using tibody at 1:1000 dilution. 5 kDa/55 kDa/60 kDa W:28 kDa/40 kDa	Western blot analysis of extracts of 293T cells using XBP1 Polyclonal Antibody at 1:1000 dilution.293T cells were treated by tunicamycin (2 μg/ml) for 8 hours. Observed-MW:35 kDa/55 kDa/60 kDa Calculated-MW:28 kDa/40 kDa	
Preparation & Storage			
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.		
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the		

Background

This gene encodes a transcription factor that regulates MHC class II genes by binding to a promoter element referred to as an X box. This gene product is a bZIP protein, which was also identified as a cellular transcription factor that binds to an enhancer in the promoter of the T cell leukemia virus type 1 promoter. It may increase expression of viral proteins by acting as the DNA binding partner of a viral transactivator. It has been found that upon accumulation of unfolded proteins in the endoplasmic reticulum (ER), the mRNA of this gene is processed to an active form by an unconventional splicing mechanism that is mediated by the endonuclease inositol-requiring enzyme 1 (IRE1). The resulting loss of 26 nt from the spliced mRNA causes a frame-shift and an isoform XBP1(S), which is the functionally active transcription factor r. The isoform encoded by the unspliced mRNA, XBP1(U), is constitutively expressed, and thought to function as a negative feedback regulator of XBP1(S), which shuts off transcription of target genes during the recovery phase of ER stress. A pseudogene of XBP1 has been identified and localized to chromosome 5.

temperature recommended.