

A Reliable Research Partner in Life Science and Medicine

NAPSA Polyclonal Antibody

catalog number: E-AB-92657

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human; Mouse; Rat

Immunogen A synthetic peptide of human NAPSA

Host Rabbit Isotype IgG


Purification Affinity purification

Buffer Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.

Recommended Dilution Applications


WB 1:1000-1:2000 IF 1:50-1:200

Data



Western blot analysis of extracts of A-549 cells using NAPSA Polyclonal Antibody at 1:500 dilution.

> Observed-MV:38 kDa Calculated-MV:45 kDa



Immunofluorescence analysis of human lung cancer cells

Immunofluorescence analysis of mouse lung cells using using NAPSA Polyclonal Antibody at dilution of 1:100 (40x NAPSA Polyclonal Antibody at dilution of 1:100 (40x lens).

Immunofluorescence analysis of rat lung cells using NAPSA

Polyclonal Antibody at dilution of 1:100 (40x lens). Blue:

DAPI for nuclear staining.

lens). Blue: DAPI for nuclear staining. Blue: DAPI for nuclear staining.

Preparation & Storage

Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles. Storage

Shipping The product is shipped with ice pack, upon receipt, store it immediately at the

temperature recommended.

Background

For Research Use Only

Rev. V1.6

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

This gene encodes a member of the peptidase A1 family of aspartic proteases. The encoded preproprotein is proteolytically processed to generate an activation peptide and the mature protease. The activation peptides of aspartic proteinases function as inhibitors of the protease active site. These peptide segments, or pro-parts, are deemed important for correct folding, targeting, and control of the activation of aspartic proteinase zymogens. The encoded protease may play a role in the proteolytic processing of pulmonary surfactant protein B in the lung and may function in protein catabolism in the renal proximal tubules. This gene has been described as a marker for lung adenocarcinoma and renal cell carcinoma.

Fax: 1-832-243-6017