

Recombinant Human NKG2A & CD94 Heterodimer Protein (His & Flag Tag)



Catalog Number:PKSH032812

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	NKG2A&CD94 Heterodimer;KLRC1&CD94 Heterodimer;CD159A&KLRD1 Heterodimer
Species	Human
Expression Host	HEK293 Cells
Sequence	Arg100-Leu233 & Ser34-Ile179
Accession	P26715&Q13241
Calculated Molecular Weight	34.4 kDa
Observed molecular weight	25-40 kDa
Tag	N-His & N-Flag

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

NKG2-A/NKG2-B Type II Integral Membrane Protein contains 1 C-type lectin domain and belongs to the killer cell lectin-like receptor family. The killer cell lectin-like receptor family is a group of transmembrane proteins preferentially expressed in NK cells. Natural killer (NK) cells are a distinct lineage of lymphocytes that mediate cytotoxic activity and secrete cytokines upon immune stimulation. CD94 (Cluster of Differentiation 94), also known as killer cell lectin-like receptor subfamily D member 1 (KLRD1), is expressed on the surface of natural killer cells in the innate immune system. CD94 Plays a role as a receptor for the recognition of MHC class I HLA-E molecules by NK cells and some cytotoxic T-cells. CD94 Can form disulfide-bonded heterodimer with NKG2 family members. The CD94/NKG2 complex, on the

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human NKG2A & CD94 Heterodimer Protein (His & Flag Tag)

Catalog Number:PKSH032812



surface of natural killer cells interacts with Human Leukocyte Antigen (HLA)-E on target cells.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017