Elabscience®

Recombinant Rhesus Macaque Angiotensin-Converting Enzyme 2/ACE-2 (C-Fc)

Catalog Number: PKSQ050120

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|---------------|--|
| Species | Rhesus Macaque |
| Source | HEK293 Cells-derived Rhesus Macaque ACE2/ACE-2 protein Gln18-Val739, with an C- |
| | terminal Fc |
| Calculated MW | 110.6 kDa |
| Observed MW | 120-140 kDa |
| Accession | ACI04564.1 |
| Bio-activity | Immobilized 2019-nCoV S Protein RBD-SD1-mFc(Cat#PKSR030476)at 2µg/ml (100 |
| | µl/well) can bind Rhesus Macaque ACE-2-Fc(Cat#PKSQ050120). The ED ₅₀ of |
| | Recombinant Rhesus Macaque ACE-2-His(Cat#PKSQ050120)is 96 ng/ml. |
| Properties | |
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Concentration | Subject to label value. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Store at $< -20^{\circ}$ C, stable for 6 months. Please minimize freeze-thaw cycles. |

packs. Upon receipt, store it immediately at < - 20°C.

Supplied as a 0.2 µm filtered solution of PBS, pH7.4.

Formulation

Shipping

<u>Da</u>ta



> 95 % as determined by reducing SDS-PAGE.



This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel

Immobilized 2019-nCoV S Protein RBD-SD1mFc(Cat#PKSR030476)at 2µg/ml (100 µl/well) can bind Rhesus Macaque ACE-2-Fc(Cat#PKSQ050120). The ED50 of Recombinant Rhesus Macaque ACE-2-His(Cat#PKSQ050120)is 96 ng/ml.

Background

Elabscience®

Angiotensin-Converting Enzyme 2 (ACE-2) is an integral membrane protein and a zinc metalloprotease of the ACE family, the ACE family includes somatic and germinal ACE. ACE-2 cleaves angiotensins I and II as a carboxypeptidase, ACE-2 converts angiotensin I to angiotensin 1-9, and angiotensin II to angiotensin 1-7. ACE-2 is also able to hydrolyze apelin-13 and dynorphin-13 with high efficiency. ACE-2 can be high expressed in testis, kidney and heart, in colon, small intestine and ovary at moderate levels. Captopril and lisinopril as the classical ACE inhibitor don't inhibit ACE-2 activity. ACE-2 may play an important role in regulating the heart function.