

Recombinant Human GAL Protein (GST Tag)

Catalog Number: PDEH100807

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	E.coli-derived Human GAL protein Gly33-Ser123, with an N-terminal GST
Calculated MW	34.9 kDa
Observed MW	35 kDa
Accession	P22466
Bio-activity	Not validated for activity

Properties

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin	< 10 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Human GAL proteins, 2 µg/lane of Recombinant Human GAL proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 35 kDa.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD180, also known as RP105, is a B-cell surface molecule belonging to the family of pathogen receptors, Toll-like receptors (TLR). CD180 has an extracellular leucine-rich repeats and a short cytoplasmic tail. CD180 / RP105 interact with an extracellular molecule named MD1 and then together form the cell surface receptor complex RP105 / MD1 which induces B-cell activation in humans and mice, leading to proliferation and up-regulation of a costimulatory molecule, B7.2 / CD86. CD180 / RP105 also has a role in LPS response because B cells lacking RP105 show hyporesponsiveness to LPS.