



A Reliable Research Partner in Life Science and Medicine

# PE/Cyanine 5.5 Anti-Human CD29 Antibody [TS2/16.2.1]

Catalog Number: E-AB-F1049I

Note: Centrifuge before opening to ensure complete recovery of vial contents.

_		4.0	
	escri	MTI.	on.
ш	COUL		UII

Reactivity Human Mouse Host

Isotype Mouse IgG1, ĸ Clone No. TS2/16.2.1

PE/Cyanine5.5 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792I] Isotype Control

PE/Cyanine 5.5 Conjugation

**Conjugation Information** PE/Cyanine5.5 is designed to be excited by the Blue (488 nm), Green (532 nm) and

yellow-green (561 nm) lasers and detected using an optical filter centered near 690 nm

(e.g., a 690/50 nm bandpass filter).

Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein

protectant.

### **Applications**

### Recommended usage

**FCM** 

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

# **Preparation & Storage**

Storage Keep as concentrated solution.

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

**Shipping** Ice bag

#### **Antigen Information**

**Alternate Names** CD29;Fibronectin receptor subunit beta;ITB1;Integrin beta-1;Itgb1;VLA-4 subunit beta

**Uniprot ID** P05556 Gene ID 3688

**Background** CD29 is a 130 kD single chain type I glycoprotein also known as integrin β1, VLA-β

> chain, or gplla. It is broadly expressed on a majority of hematopoietic and nonhematopoietic cells, including leukocytes (although at low level on granulocytes), platelets, fibroblasts, endothelial cells, epithelial cells, and mast cells. CD29 is a member of the integrin family. It is non-covalently associated with integrin α1-α6 chains to form VLA-1 to VLA-6 molecules, respectively. Integrins, which include CD29, bind to several cell surface (e.g. VCAM-1, MadCAM-1) and extracellular matrix molecules. CD29 acts as a fibronectin receptor and is involved in a variety of cell-cell and cell-

matrix interactions.

# For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Web:www.elabscience.com

Email:techsupport@elabscience.com