

Recombinant Human GRK2/ADRBK1 Protein (His & GST Tag)

Catalog Number: PKSH031009

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	Baculovirus-Insect Cells-derived Human GRK2/ADRBK1 protein Met 1-Leu 689, with an N-terminal His & GST
Calculated MW	107 kDa
Observed MW	110 kDa
Accession	NP_001610.2
Bio-activity	Not validated for activity

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 50mM Tris, 500mM NaCl, 0.5mM GSH, pH 8.0 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V3.6

G-protein coupled receptor kinase 2 (GRK2), also referred as Adrenergic, beta, receptor kinase 1 (ADRBK1), is a ubiquitous member of the Gprotein-coupled receptor kinase (GRK) family that appears to play a central, integrative role in signal transduction cascades. GRK2 can phosphorylate a growing number of non-GPCR substrates and associate with a variety of proteins related to signal transduction, thus suggesting that this kinase could also have diverse 'effector' functions. GRK2 has been reported to interact with a variety of signal transduction proteins related to cell migration such as MEK, Akt, PI3Kgamma or GIT. Interestingly, the levels of expression and activity of this kinase are altered in a number of inflammatory disorders (as rheumatoid arthritis or multiple sclerosis), thus suggesting that GRK2 may play an important role in the onset or development of these pathologies. The important physiological function of GRK2 as a modulator of the efficacy of GPCR signal transduction systems is exemplified by its relevance in cardiovascular physiopathology as well as by its emerging role in the regulation of chemokine receptors. Besides its canonical role in the modulation of the signalling mediated by many Gprotein-coupled receptors (GPCR), this protein can display a very complex network of functional interactions with a variety of signal transduction partners, in a stimulus, cell type, or context-specific way.