

## Recombinant Human RELT/TNFRSF19L Protein (His Tag)

Catalog Number: PKSH031543

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Human
<b>Source</b>	HEK293 Cells-derived Human RELT/TNFRSF19L protein Met 1-Ala 160, with an C-terminal His
<b>Calculated MW</b>	15.7 kDa
<b>Accession</b>	NP_116260.2
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

#### For Research Use Only

Receptor expressed in lymphoid tissues (RELT); also known as tumor necrosis factor receptor superfamily, member 19-like (TNFRSF19L); is a member of the TNF-receptor superfamily. This receptor is especially abundant in hematologic tissues. It has been shown to activate the NF-kappaB pathway and selectively bind TNF receptor-associated factor 1. RELT/TNFRSF19L is capable of stimulating T-cell proliferation in the presence of CD3 signaling; which suggests its regulatory role in immune response. RELT/TNFRSF19L is a type I transmembrane glycoprotein with a cysteine-rich extracellular domain; possessing significant homology to other members of the TNFR superfamily; especially TNFRSF19; DR3; OX40; and LTbeta receptor. RELT/TNFRSF19L is able to activate the NF-kappaB pathway and selectively binds tumor necrosis factor receptor-associated factor 1. RELT/TNFRSF19L is able to activate the NF-&kappa;B pathway and selectively binds tumor necrosis factor receptor-associated factor 1. Although the soluble form of RELT fusion protein does not inhibit the one-way mixed lymphocyte reaction; immobilized RELT/TNFRSF19L is capable of costimulating T-cell proliferation in the presence of CD3 signaling.

## For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017