

Recombinant Human Frizzled-8 (C-6His-Avi) Biotinylated

Catalog Number: PKSH034044

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	HEK293 Cells-derived Human Frizzled-8 protein Ala28-Pro172, with an C-terminal His & Avi
Calculated MW	19.1 kDa
Observed MW	25-35 kDa
Accession	Q9H461
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 °C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization.
Reconstitution	Please refer to the specific buffer information in the printed manual. Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Frizzled-8 is one of at least ten seven-transmembrane (7TM) glycoproteins of the Frizzled family of Wnt receptors. Frizzled proteins are thought to be G-protein-coupled. Wnt engagement, with low density lipoprotein receptor-related proteins LRP-5 or LRP-6 acting as co-receptors, stabilizes beta -catenin and promotes gene transcription that is important in development and tissue maintenance. Component of the Wnt-Fzd-LRP5-LRP6 complex that triggers beta-catenin signaling through inducing aggregation of receptor-ligand complexes into ribosome-sized signalosomes. The beta-catenin canonical signaling pathway leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes. These ligands bind the extracellular CRD of Frizzled-8, blocking Wnt binding. The recombinant Frizzled-8 CRD has also been used to block Wnt signaling and inhibit growth of teratocarcinomas.