

Recombinant Human AIFM1 Protein (His Tag)

Catalog Number: PKSH032089

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Species	Human
Source	E.coli-derived Human AIFM1 protein Glu121-Asp613, with an N-terminal His
Calculated MW	56.2 kDa
Observed MW	68 kDa
Accession	O95831
Bio-activity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Concentration	Subject to label value.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < - 20°C.
Formulation	Supplied as a 0.2 µm filtered solution of PBS, 50% Glycerol, 2mM EDTA, 0.5M Arginine, 5% Trehalose, pH 7.4.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Apoptosis-Inducing Factor 1, Mitochondrial (AIFM1) is a flavoprotein essential for nuclear disassembly in apoptotic cells that is found in the mitochondrial intermembrane space in healthy cells. During apoptosis, it is translocated from the mitochondria to the nucleus to function as a proapoptotic factor in a caspase-independent pathway, while in normal mitochondria, it functions as an antiapoptotic factor via its oxidoreductase activity. The soluble form (AIFsol) found in the nucleus induces parthanatos i.e., caspase-independent fragmentation of chromosomal DNA. AIFM1 interacts with EIF3G, and thereby inhibits the EIF3 machinery and protein synthesis, and activates caspase-7 to amplify apoptosis. It binds to DNA in a sequence-independent manner and plays a critical role in caspase-independent, pyknotic cell death in hydrogen peroxide-exposed cells.

For Research Use Only