

# Recombinant Human Rad51 protein (His tag)

Catalog Number:PDEH100341



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Description

|                                    |  |
|------------------------------------|--|
| <b>Synonyms</b>                    | DNA repair protein RAD51 homolog 1;RAD51; HsRAD51;hRAD51 |
| <b>Species</b>                     | Human  |
| <b>Expression Host</b>             | E.coli   |
| <b>Sequence</b>                    | Ala 2-Asp 339  |
| <b>Accession</b>                   | Q06609   |
| <b>Calculated Molecular Weight</b> | 37.1 kDa   |
| <b>Observed molecular weight</b>   | 38 kDa   |
| <b>Tag</b>                         | N-His  |

## Properties

|                       |   |
|-----------------------|---|
| <b>Purity</b>         | > 95 % as determined by reducing SDS-PAGE.  |
| <b>Endotoxin</b>      | Please contact us for more information.   |
| <b>Storage</b>        | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| <b>Shipping</b>       | This product is provided as lyophilized powder which is shipped with ice packs.   |
| <b>Formulation</b>    | Lyophilized from sterile PBS, pH 7.4.<br>Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.<br>Please refer to the specific buffer information in the printed manual.           |
| <b>Reconstitution</b> | Please refer to the printed manual for detailed information.  |

## Background

Plays an important role in homologous strand exchange, a key step in DNA repair through homologous recombination. Binds to single and double-stranded DNA and exhibits DNA-dependent ATPase activity. Catalyzes the recognition of homology and strand exchange between homologous DNA partners to form a joint molecule between a processed DNA break and the repair template. Binds to single-stranded DNA in an ATP-dependent manner to form nucleoprotein filaments which are essential for the homology search and strand exchange (PubMed:26681308). Part of a PALB2-scaffolded HR complex containing BRCA2 and RAD51C and which is thought to play a role in DNA repair by HR. Plays a role in regulating mitochondrial DNA copy number under conditions of oxidative stress in the presence of RAD51C and XRCC3.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017