Recombinant Mouse IFNα2 Protein(His Tag)

Catalog Number: PDMM100239

Note: *Centrifuge before opening to ensure complete recovery of vial contents.*

Description

Species	Mouse
Source	Mammalian-derived Mouse IFNα2 protein Cys24-Glu190, with an C-teminal His
Calculated MW	18.2 kDa
Observed MW	18-20 kDa
Accession	P01573
Bio-activity	Not validated for activity

Properties

Purity	> 90% as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU/mg of the protein as determined by the LAL method
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80 ℃. Reconstituted protein solution can be stored at 4-8℃ for 2-7 days. Aliquots of reconstituted samples are stable at < -20℃ for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis.

Data



SDS-PAGE analysis of Mouse IFNα2 proteins, 2μg/lane of Recombinant Mouse IFNα2 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 18-20 kDa

Background

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web:www.elabscience.com
Email:techsupport@elabscience.com

IFNA2 (Interferon Alpha 2) is a Protein Coding gene. This gene is a member of the alpha interferon gene cluster on chromosome 9. The encoded protein is a cytokine produced in response to viral infection. Type I Interferons (IFNs) are well-known cytokines that exert antiviral activity, antitumor activity, and immunomodulatory effects. Interferon tau (IFN T), a type I IFN similar to alpha IFNs (IFNA), is the pregnancy recognition signal produced by the ruminant conceptus. Among the IFN-α genes, a total of 28 different sequence variants have been described. The three principal subtypes of IFNα-2 are designated α-2a, α-2b, and α-2c. IFNα-2b is being the predominant allele while IFNα-2a is less predominant and IFNα-2c only a minor allelic variant.