

A Reliable Research Partner in Life Science and Medicine

Recombinant PIGR Monoclonal Antibody

catalog number: AN300080P

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human


Immunogen Recombinant Human PIGR protein

Host Rabbit Isotype IgG 12G8 Clone **Purification** Protein A


Buffer 0.2 µm filtered solution in PBS

Applications	Recommended Dilution	
IHC-P	1:50-1:200	
ICC/IF	1:20-1:100	

Data




Immunofluorescence analysis of PIGR in HepG2 cells. Cells were fixed with 4% PFA, blocked with 10% serum, and incubated with rabbit anti-Y PIGR Monoclonal Antibody (dilution ratio 1:60) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI for nuclear staining (blue). Positive staining was localized to



Immunohistochemistry of paraffin-embedded human breast carcinoma using PIGR Monoclonal Antibody at dilution of

1:60.



Immunohistochemistry of paraffin-embedded human liver cirrhosis using PIGR Monoclonal Antibody at dilution of 1:60.

Preparation & Storage

Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.

For Research Use Only

Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Email:techsupport@elabscience.com



Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Shipping Ice bag

Background

The human polymeric immunoglobulin receptor (pIgR; also known as membrane secretory component) is an approximately 100 kDa type I transmembrane glycoprotein that is synthesized as a 764 amino acid (aa) precursor. It includes a signal sequence (aa 1-18), an extracellular region (aa 19-638), a transmembrane segment (aa 639-661), and a cytoplasmic domain (aa 662-764). The extracellular region consists of five Ig-like domains and a sixth non-Ig domain that connects to the membrane region. pIgR is expressed on secretory epithelial cells of exocrine tissues. Immunoglobulin isotypes consist of two heavy (H) and two light (L) chains. F

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Email:techsupport@elabscience.com Web:www.elabscience.com