

Recombinant MAP2 Monoclonal Antibody

catalog number: **AN300528P**

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Mouse
Immunogen	Recombinant Mouse MAP2 protein
Host	Rabbit
Isotype	IgG
Clone	7A2
Purification	Protein A
Buffer	0.2 µm filtered solution in PBS

Applications


Recommended Dilution

IHC-P	1:500-1:2000
--------------	--------------


Data



Immunohistochemistry of paraffin-embedded mouse spleen using MAP2 Monoclonal Antibody at dilution of 1:1000.



Immunohistochemistry of paraffin-embedded mouse kidney using MAP2 Monoclonal Antibody at dilution of 1:1000.



Immunohistochemistry of paraffin-embedded mouse intestine using MAP2 Monoclonal Antibody at dilution of 1:1000.

Preparation & Storage

Storage	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Shipping	Ice bag

Background

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.0

METAP2 (Methionine aminopeptidase 2), also known as MAP2 is a protein that belongs to the peptidase M24A family. MAP2 binds 2 cobalt or manganese ions and contains approximately 12 O-linked N-acetylglucosamine (GlcNAc) residues. It is found in all organisms and is especially important because of its critical role in tissue repair and protein degradation. The catalytic activity of human MAP2 toward Met-Val peptides is consistently two orders of magnitude higher than that of METAP1, suggesting that it is responsible for processing proteins containing N-terminal Met-Val and Met-Thr sequences in vivo. This protein functions both by protecting the alpha subunit of eukaryotic initiation factor 2 from inhibitory phosphorylation and by removing the amino-terminal methionine residue from nascent protein. MAP2 protects eukaryotic initiation factor EIF2S1 from translation-inhibiting phosphorylation by inhibitory kinases such as EIF2AK2/PKR and EIF2AK1/HCR. It also plays a critical role in the regulation of protein synthesis.