

Recombinant TRIAD1 Monoclonal Antibody

catalog number: **AN302043L**

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human;
Immunogen	Peptide. This information is proprietary to PTMab.
Host	Rabbit
Isotype	IgG, κ
Clone	A763
Purification	Protein A purified
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protein protectant.

Applications Recommended Dilution

WB	1:1000
-----------	--------

Data



Western Blot with TRIAD1 Monoclonal Antibody at dilution of 1:1000. Lane 1: Jurkat, Lane 2: 293T, Lane 3: Ramos

Observed-MW:58 kDa

Calculated-MW:58 kDa

Preparation & Storage

Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	Ice bag

Background

The E3 ubiquitin-protein ligase ARIH2 (TRIAD1) is an Ariadne subfamily ligase involved in the polyubiquitination of proteins designated for proteasomal degradation. The TRIAD1 nuclear protein contains an amino-terminal acidic region, a pair of RING fingers, two carboxyl terminal coiled coil domains and a novel C6HC DRIL/IBR domain located between the RING fingers. Together, the paired RING fingers and DRIL/IBR domain form a highly conserved TRIAD (two RING fingers and DRIL) domain. Research studies suggest that TRIAD1 mediates both Lys48 and Lys63 protein polyubiquitination and acts as a negative regulator of myelopoiesis. TRIAD1 ubiquitin ligase inhibits myeloid cell proliferation by mediating protein ubiquitination through the ubiquitin-conjugating enzymes UbcH7 and UbcH13. Experimental deletion of TRIAD1 in mice has a lethal effect, leading to death at the embryonic stage or later due to a severe, multi-organ inflammatory response. Results indicate that TRIAD1 binds I κ B β in dendritic cells and promotes the degradation of the NF- κ B inhibitor.

For Research Use Only

Toll-free: 1-888-852-8623
Web: www.elabscience.com

Tel: 1-832-243-6086
Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Rev. V1.0