# **Elabscience**<sup>®</sup>

## **PPM1D** Polyclonal Antibody

### catalog number: E-AB-52172


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Reactivity	Human;Mouse	
Immunogen	Fusion protein of human PPM1D	
Host	Rabbit	
Isotype	IgG	
Purification	Antigen affinity purification	
Buffer	Phosphate buffered solution, pH 7.4, containing 0.05% stabilizer and 50% glycerol.	

Applications	Recommended Dilution
IHC	1:50-1:300

#### Data





Immunohistochemistry of paraffin-embedded Human brain Immunohistochemistry of paraffin-embedded Human ovarian tissue using PPM1D Polyclonal Antibody at dilution of 1:60(×200)

cancer tissue using PPM1D Polyclonal Antibody at dilution of 1:60(×200)

Preparation & Storage	
Storage	Store at -20°C Valid for 12 months. Avoid freeze / thaw cycles.
Shipping	The product is shipped with ice pack, upon receipt, store it immediately at the temperature recommended.

#### Background

The protein encoded by this gene is a member of the PP2C family of Ser/Thr protein phosphatases. PP2C family members are known to be negative regulators of cell stress response pathways. The expression of this gene is induced in a p53dependent manner in response to various environmental stresses. While being induced by tumor suppressor protein TP53/p53, this phosphatase negatively regulates the activity of p38 MAP kinase, MAPK/p38, through which it reduces the phosphorylation of p53, and in turn suppresses p53-mediated transcription and apoptosis. This phosphatase thus mediates a feedback regulation of p38-p53 signaling that contributes to growth inhibition and the suppression of stress induced apoptosis. This gene is located in a chromosomal region known to be amplified in breast cancer. The amplification of this gene has been detected in both breast cancer cell line and primary breast tumors, which suggests a role of this gene in cancer development.