

## Recombinant Mouse VDR/NR1I1 Protein (His Tag)

**Catalog Number:** PKSM040354

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Mouse
<b>Source</b>	Baculovirus-Insect Cells-derived Mouse VDR/NR1I1 protein Met1-Ser422, with an C-terminal His
<b>Calculated MW</b>	49.2 kDa
<b>Observed MW</b>	55 kDa
<b>Accession</b>	P48281
<b>Bio-activity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 86 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 8.0, 10% glycerol Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 86 % as determined by reducing SDS-PAGE.

### Background

#### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

VDR (vitamin D(1,25- dihydroxyvitamin D3)receptor), also known as NR1H1, belongs to the NR1H1 family, NR1 subfamily. It is composed of three domains: a modulating N-terminal domain, a DNA-binding domain and a C-terminal ligand-binding domain. Vitamin D receptors (VDRs) are members of the NR1H1 family, which also includes pregnane X (PXR) and constitutive androstane (CAR) receptors, that form heterodimers with members of the retinoid X receptor family. VDRs repress expression of 1 $\alpha$ -hydroxylase (the proximal activator of 1,25(OH) $_2$ D $_3$ ) and induce expression of the 1,25(OH) $_2$ D $_3$  inactivating enzyme CYP24. Also, it has recently been identified as an additional bile acid receptor alongside FXR and may function to protect gut against the toxic and carcinogenic effects of these endobiotics. VDR is expressed in the intestine, thyroid and kidney and has a vital role in calcium homeostasis. It is the nuclear hormone receptor, also called transcription factor that mediates the action of vitamin D $_3$ . Inherited mutations in the VDR gene leads to rickets.

## For Research Use Only

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017