PERK Polyclonal Antibody

Catalog Number: E-AB-16703



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human, Mouse, Rat

Immunogen Synthetic peptide of human EIF2AK3

Host Rabbit
Isotype IgG

Purification Affinity purification
Conjugation Unconjugated

Formulation PBS with 0.05% sodium azide and 50% glycerol, PH7.4

Applications Recommended Dilution

IHC 1:50-1:200

Data



Immunohistochemistry of paraffin-embedded Human prostate cancer tissue using PERK Polyclonal Antibody at dilution 1:50



Immunohistochemistry of paraffin-embedded Human brain tissue using PERK Polyclonal Antibody at dilution 1:50

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

The protein encoded by this gene phosphorylates the alpha subunit of eukaryotic translation-initiation factor 2 (EIF2), leading to its inactivation, and thus to a rapid reduction of translational initiation and repression of global protein synthesis. It is a type I membrane protein located in the endoplasmic reticulum (ER), where it is induced by ER stress caused by malfolded proteins. Mutations in this gene are associated with Wolcott-Rallison syndrome.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com