

Noggin/NOG (N-8His-Flag), Human, Recombinant

Cat. No. : PCK211

General Information

Synonyms	Noggin;NOG
Species	Human
Expression host	Human Cells
Sequence	Gln28-Cys232
Accession	Q13253
Mol mass	23 kDa
Expiration date	12 months
Bio activity	Measured by its ability to inhibit BMP-4 induced alkaline phosphatase production by ATDC5 mouse chondrogenic cells. The ED50 for this effect is 10-100 ng/mL in the presence of 200 ng/mL of Recombinant Human BMP-4.


Product feature

Purity	> 95% as determined by reducing SDS-PAGE.
Endotoxin (EU/μg)	< 0.1
Storage	Lyophilized protein should be stored at -5~-20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -5~-20°C for 3 months.
Shipping	Ice bag
Formulation	Lyophilized from a 0.2 μm filtered solution of 20 mM PB, 500 mM NaCl, 2 mM EDTA, pH 7.4
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μg/mL. Dissolve the lyophilized protein in sterile water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Background

Noggin is a secreted homodimeric glyco Protein that is an antagonist of bone morphogenetic Proteins (BMPs). Mature Human Noggin contains an N-terminal acidic region, a central basic heparin-binding segment and a C-terminal cysteine-knot structure. Noggin is very highly conserved among vertebrates, such that mature human Noggin shares 99%, 99%, 98%, 97% and 89% aa sequence identity with mouse, rat bovine, equine and chicken Noggin, respectively. Secreted Noggin probably remains close to the cell surface due to its binding of heparin-containing proteoglycans. Noggin binds some BMPs such as BMP4 with high affinity and others such as BMP7 with lower affinity. It antagonizes BMP bioactivities by blocking epitopes on BMPs that are needed for binding to both type I and type II Receptors. Noggin is expressed in defined areas of the adult central nervous system and peripheral tissues such as lung, skeletal muscle and skin. During culture of human embryonic stem cells (hESC) or neural stem cells under certain conditions, addition of Noggin to antagonize BMP activity may allow stem cells to proliferate while maintaining their undifferentiated state, or alternatively, to differentiate into dopaminergic neurons.

SDS-PAGE



Price¹
by Elabscience

Pricella[®]
by Elabscience

Pricella[®]
by Elabscience

Pricella[®]
by Elabscience