

A Reliable Research Partner in Life Science and Medicine

## Recombinant Mouse IGFBP-2 / IGFBP2 Protein (His Tag)

Catalog Number: PDEM100304

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### Description

**Species** Mouse

Source E.coli-derived Mouse IGFBP-2 protein Glu35-Gln305, with an N-terminal His

**Calculated MW** 29.7 kDa Observed MW 39 kDa P47877 Accession

Not validated for activity **Bio-activity** 

#### **Properties**

> 95% as determined by reducing SDS-PAGE. **Purity** 

**Endotoxin** < 10 EU/mg of the protein as determined by the LAL method

Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -Storage

80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.


Shipping This product is provided as lyophilized powder which is shipped with ice packs. **Formulation** 

Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5%

Reconstitution It is recommended that sterile water be added to the vial to prepare a stock solution

of 0.5 mg/mL. Concentration is measured by UV-Vis.

### Data



SDS-PAGE analysis of Mouse IGFBP-2 / IGFBP2 proteins, 2 μg/lane of Recombinant Mouse IGFBP-2 / IGFBP2 proteins was resolved with SDS-PAGE under reducing conditions, showing bands at 39 kDa.

#### **Background**

For Research Use Only

Toll-free: 1-888-852-8623 Fax: 1-832-243-6017 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com

# Elabscience®

#### Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

IGFBP-2, also known as IGFBP2, is a insulin-like growth factor-binding protein (IGFBP). IGFBPs prolong the half-life of the IGFs, control bioavailability, activity, and distribution of insulin-like growth factor (IGF) through high-affinity IGFBP/ IGF complexes. Six high-affinity IGF-binding proteins (IGFBP-1 to-6) have been identified. The six IGFBPs are structurally related but encoded by distinct genes. IGFBPs have a high affinity for IGFs. Some members of the IGFBP family have been consistently shown to inhibit IGF actions by preventing them from gaining access to the IGF receptors, while others potentiate IGF actions by facilitating the ligand-receptor interaction. IGFBP-2 is overexpressed in many malignancies and is often correlated with an increasingly malignant status of the tumor, pointing to a potential involvement of IGFBP-2 in tumorigenesis. It contains 1 IGFBP N-terminal domain and 1 thyroglobulin type-1 domain. It inhibits IGF-mediated growth and developmental rates.

For Research Use Only

 Toll-free: 1-888-852-8623
 Tel: 1-832-243-6086
 Fax: 1-832-243-6017

 Web: www.elabscience.com
 Email: techsupport@elabscience.com