

## Recombinant Mouse CD155/PVR Protein (aa 1-345, His Tag)

Catalog Number: PKSM040773

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Species</b>	Mouse
<b>Source</b>	HEK293 Cells-derived Mouse CD155/PVR protein Met 1-Arg 345, with an C-terminal His
<b>Calculated MW</b>	28.0 kDa
<b>Observed MW</b>	60-65 kDa
<b>Accession</b>	NP_081790.1
<b>Bio-activity</b>	Measured by its ability to bind recombinant mouse CD226/DNAM-1. Immobilized recombinant mouse CD155/PVR at 1 µg/ml (100 µl/well) can bind recombinant mouse CD226/DNAM-1 with a linear range of 0.78-100 ng/ml.

### Properties

<b>Purity</b>	> 97 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



Measured by its ability to bind recombinant mouse CD226/DNAM-1. Immobilized recombinant mouse CD155/PVR at 1 µg/ml (100 µl/well) can bind recombinant mouse CD226/DNAM-1 with a linear range of 0.78-100 ng/ml.

### Background

#### For Research Use Only

Toll-free: 1-888-852-8623  
Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086  
Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

CD155, commonly known as PVR (poliovirus receptor) and Necl-5 (nectin-like molecule-5), is a type I transmembrane single-span glycoprotein, and belongs to the nectins and nectin-like (Necl) subfamily. CD155 was originally identified based on its ability to mediate the cell attachment and entry of poliovirus (PV), an etiologic agent of the central nervous system disease poliomyelitis. The normal cellular function is in the establishment of intercellular adherens junctions between epithelial cells. CD155 may assist in an efficient humoral immune response generated within the intestinal immune system. It has been demonstrated that CD155 can be recognized and bond by DNAM-1 and CD96 which promote the adhesion, migration and NK-cell killing, and thus efficiently prime cell-mediated tumor-specific immunity.

## For Research Use Only

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017