A Reliable Research Partner in Life Science and Medicine

# Recombinant Human ILT6/LILRA3 Protein (Fc Tag)

Catalog Number: PKSH030602

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### Description

Species Human

Source HEK293 Cells-derived Human ILT6/LILRA3 protein Met 1-Glu439, with an C-terminal

hFc

Calculated MW 72.0 kDa
Observed MW 102 kDa
Accession AAH28208.1

**Bio-activity** Not validated for activity

#### **Properties**

**Purity** > 80 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80

°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, pH 7.4


Normally 5% - 8% trehalose, mannitol and 0.01% Tween 80 are added as protectants

before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 80 % as determined by reducing SDS-PAGE.

#### **Background**

ILT6, also known as LILRA3, belongs to the ILT family. In human, the ILT gene family includes up to 11 members. The extracellular portion of all members includes at least two and usually four immuno-globulin domains. ILT-2 through 5 are all inhibitory members having variable numbers of cytoplasmic ITIM domains. ILT6 lacks a transmembrane domain. The function of ILT6 is currently unknown. however it is highly homologous to other LILR genes, and can bind human leukocyte antigen (HLA) class I. Therefore, if secreted, the ILT6 might impair interactions of membrane-bound LILRs ( such as LILRB1, an inhibitory receptor expressed on effector and memory CD8 T cells) with their HLA ligands, thus modulating immune reactions and influencing susceptibility to disease.

## For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017