

A Reliable Research Partner in Life Science and Medicine

APC Anti-Human CD14 Antibody[M5E2]

Catalog Number: E-AB-F1209E

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human Mouse Host

Isotype Mouse IgG2a, ĸ

Clone No. M5E2

APC Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802E] Isotype Control

APC Conjugation

Conjugation Information APC is designed to be excited by the Red (627-640 nm) laser and detected using an

optical filter centered near 660 nm (e.g., a 660/20 nm bandpass filter).


Storage Buffer Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer.

Applications Recommended usage

FCM

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 μL staining volume or per 100 μL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Data



Human pheripheral blood monocytes are stained with APC Anti-Human CD14 Antibody (filled gray histogram). Unstained pheripheral blood monocytes (blank black histogram) are used as control.

Preparation & Storage

Keep as concentrated solution. **Storage**

This product can be stored at 2-8°C for 12 months. Please protected from prolonged

exposure to light and do not freeze.

Shipping Ice bag

Antigen Information

Alternate Names CD14;Monocyte differentiation antigen CD14;Myeloid cell-specific leucine-rich

glycoprotein

Uniprot ID P08571 Gene ID 929

For Research Use Only

Tel: 1-832-243-6086 Fax: 1-832-243-6017 Toll-free: 1-888-852-8623 Email:techsupport@elabscience.com

Web:www.elabscience.com

Elabscience[®]

Elabscience Bionovation Inc.

A Reliable Research Partner in Life Science and Medicine

Background

CD14 is a 53-55 kD glycosylphosphatidylinositol (GPI)-linked membrane glycoprotein also known as LPS receptor. CD14 is expressed at high levels on monocytes and macrophages, and at lower levels on granulocytes. Some dendritic cell populations such as interfollicular dendritic cells, reticular dendritic cells, and Langerhans cells have also been reported to express CD14. As a high-affinity receptor for LPS, CD14 is involved in the clearance of gram-negative pathogens, and in the upregulation of adhesion molecules and expression of cytokines in monocytes and neutrophils.

Fax: 1-832-243-6017